
MARTIN–LUTHER–UNIVERSITÄT
HALLE–WITTENBERG
INSTITUT FÜR MATHEMATIK



**The relationship between multicriteria
robustness concepts and set-valued
optimization**

**J. Ide, E. Köbis, D. Kuroiwa, A. Schöbel and
Chr. Tammer**

Report No. 04 (2013)

Editors:

Professors of the Institute for Mathematics, Martin-Luther-University Halle-Wittenberg.

Electronic version: see <http://www2.mathematik.uni-halle.de/institut/reports/>

The relationship between multicriteria robustness concepts and set-valued optimization

J. Ide, E. Köbis, D. Kuroiwa, A. Schöbel and
Chr. Tammer

Report No. 04 (2013)

Jonas Ide
Anita Schöbel
Institut für Numerische und Angewandte Mathematik Lotzestraße 16-18
D-37083 Göttingen
Email: j.ide@math.uni-goettingen.de
schoebel@math.uni-goettingen.de

Daishi Kuroiwa
Department of Mathematics
Interdisciplinary Faculty of Science and Engineering Shimane University
Email: kuroiwa@math.shimane-u.ac.jp

Elisabeth Köbis
Christiane Tammer
Martin-Luther-Universität Halle-Wittenberg
Naturwissenschaftliche Fakultät II
Institut für Mathematik
Theodor-Lieser-Str. 5
D-06120 Halle/Saale, Germany
Email: elisabeth.koebis@mathematik.uni-halle.de
christiane.tammer@mathematik.uni-halle.de

The relationship between multicriteria robustness concepts and set-valued optimization

Jonas Ide ¹, Elisabeth Köbis, Daishi Kuroiwa, Anita Schöbel and Christiane Tammer

Dedicated to Prof. W. Takahashi on the occasion of his 70th birthday

Abstract

In this paper, we discuss the connection between concepts of robustness for multi-objective optimization problems and set order relations. We extend some of the existing concepts to general spaces and cones using set relations. Furthermore, we derive new concepts of robustness for multi-objective optimization problems. We point out that robust multi-objective optimization can be interpreted as an application of set-valued optimization. Furthermore, we develop new algorithms for solving uncertain multi-objective optimization problems. These algorithms can be used in order to solve a special class of set-valued optimization problems.

Keywords. Robust optimization, multi-objective optimization, scalarization, vectorization, set-valued optimization

1 Introduction

Dealing with uncertainty in multi-objective optimization problems is very important in many applications. On the one hand, most real world optimization problems are contaminated with uncertain data, especially traffic optimization problems, scheduling problems, portfolio optimization, network flow and network design problems. On the other hand, many real world optimization problems require the minimization of multiple conflicting objectives (see [28]), e.g. the maximization of the expected return versus the minimization of risk in portfolio optimization, the minimization of production time versus the minimization of the cost of manufacturing equipment, or the maximization of tumor control versus the minimization of normal tissue complication in radiotherapy treatment design.

For an optimization problem contaminated with uncertain data it is typical that at the time it is solved these data are not completely known. It is very important to estimate the effects of this uncertainty and so it is necessary to evaluate how sensitive an optimal solution is to perturbations of the input data. One way to deal with this question is *sensitivity analysis* (for an overview see [25]). Sensitivity analysis is an a posteriori approach and provides ranges for input data within which a solution remains feasible or optimal. It does not, however, provide a course of action for changing a solution should the perturbation be outside this range. In contrast, *stochastic programming* (see e.g. Birge and Louveaux [5] for an introduction) and *robust optimization* (see e.g. [16, 3] for an overview) take the uncertainty into account during the optimization process. While stochastic programming assumes some knowledge about the probability distribution of the uncertain data and the objective usually is to find a solution that is feasible with a certain probability and that optimizes the expected value of some objective function, robust optimization hedges against the worst case. Hence robust optimization does not require any probabilistic information. Depending from the concrete application one can decide whether robust or stochastic optimization is the more appropriate way of dealing with uncertainty.

Robust optimization is usually applied to problems where a solution is required which hedges against all possible scenarios. For example, the emergency department with landing place for rescue helicopters in a ski resort should be chosen in such a way that the flight time to all ski slopes in the resort that are to be protected is minimized in the worst case, even though flight times are uncertain due to unknown weather conditions. Similarly, if an aircraft-schedule of an airline is to be determined, one would want to be able to provide service to as many passengers as possible in a cost-effective manner, even though the exact number of passengers is not known at the time the schedule is fixed.

Generally, in the concept of robustness it is not assumed that all data are known, but one allows different scenarios for the input parameters and looks for a solution that works well in every uncertain scenario.

¹Supported by DFG RTG 1703 “Resource Efficiency in Interorganizational Networks”

Unfortunately, at the time the uncertain optimization problem has to be solved, it is not known which scenario is going to be realized. Therefore, a definition of a “good” (or robust against the perturbations in the uncertain parameter) solution is necessary.

Robust optimization is a growing field of research, we refer to Ben-Tal, L. El Ghaoui, Nemirovski [3], Kouvelis and Yu [16] for an overview of results and applications for the most prominent concepts. Several other concepts of robustness were introduced more recently, e.g. the concept of light robustness by Fischetti and Monaci [11] or the concept of recovery-robustness in Liebchen et al. [23], for a unified approach, see [15]. A scenario-based approach is suggested in Goerigk and Schöbel [12]. In all these approaches, the uncertain optimization problem is replaced by a deterministic version, called the *robust counterpart* of the uncertain problem.

One of the most common approaches is the concept of minmax robustness, introduced by Soyster [26] and studied e.g. by Ben-Tal and Nemirovski [4]. Here, a solution is said to be robust, if it minimizes the worst case of the objective function over all scenarios. We do not go into detail here as for this paper we mostly consider concepts of robustness for multi-objective optimization problems.

Now, if we consider the objective function in the problem definition to be not a single-objective, but a multi-objective function, the concepts of robustness do not apply naturally anymore. The problem obviously is that there is no total order on \mathbb{R}^k and the robustness concepts for uncertain single-objective optimization problems rely on the total order of \mathbb{R} . Therefore, new definitions of what is seen as a robust solution to an uncertain multi-objective optimization problem are necessary.

The first approach to handling uncertainty for multi-objective optimization problems was presented by Deb and Gupta [7] who extended the concept Branke [6] introduced for single-objective functions. Here each objective function is replaced by their mean function and an efficient solution to the resulting multi-objective optimization problem is called a robust solution. The authors also presented a second definition where the uncertainty is modeled into the constraints which restrict the variation of the original objective functions to their means. Barrico and Antunes [2] extended the concept of Deb and Gupta and introduced the degree of robustness as a measure how much a predefined neighborhood of the solution considered can be extended without containing solutions whose function values are too bad. An overview of the existing concepts of robustness for multi-objective optimization problems can be found in [27] and [32].

A first approach to extending the concept of minmax robustness to multi-objective optimization was presented by Kuroiwa and Lee [21]. Here, the worst case in each component is calculated separately, and an efficient solution to the problem of minimizing the vector of worst cases is then called a robust solution to the original problem. This definition has been extended by Ehrgott, et al. [8], where the authors replace the objective function by a set-valued objective function. Furthermore, the authors present solution algorithms for calculating minmax robust efficient solutions, one of which is closely connected to the concept of robustness presented by Kuroiwa and Lee [21]. Furthermore, the authors in [21] present solution concepts for obtaining robust points of uncertain multi-objective optimization and study optimality conditions for the special case of convex objective functions in [22].

Set-valued optimization on the other hand deals with the problem of minimizing a function where the image of a point is in fact a set. Minimizing a set is not totally intuitive since on a power set there is no total order as well as on \mathbb{R}^k . Therefore, a definition of what can be seen as an optimal solution to minimizing a set-valued objective function is necessary. In order to compare sets, several preorders have been introduced (see e.g. [10, 17, 20, 24, 33]). With these preorders it is then possible to formulate set-valued optimization problems related to robustness for multi-objective optimization problems, especially, we show that the concept of minmax robust efficiency (see [8]) is closely connected to a certain set order relation, introduced by Kuroiwa [17, 20], namely the *upper-type set-relation*. We derive our results in general spaces using arguments from nonlinear and convex analysis (see Takahashi [29, 30]), for methods from numerical analysis in general spaces, see e.g. Aoyama, Kohsaka, Takahashi [1], Takahashi [31].

Replacing the set order relation implicitly used in the definition of minmax robust efficiency, Ide and Köbis [14] presented various other concepts of robustness for multi-objective optimization, derived by replacing the upper-type set-relation with another set ordering from the literature. The authors then presented algorithms to calculate the resulting robust efficient solutions.

Now, this paper is structured as follows: After fixing the notation and recalling the definitions of set order relations in Section 2, in Section 3 we introduce several concepts of robustness for multi-objective optimization problems based on set order relations. We show some characterizations for robust solutions in the sense of set-valued optimization that are important for deriving solution procedures using the ideas given in [8]. A lot of the results presented in [8] can be extended to our general setting. Using this information, we extend the algorithms presented in [8] to concepts for robustness and then use these algorithms in order

to solve a certain class of set-valued optimization problems. We conclude the paper with some final remarks and an outlook to future research.

2 Preliminaries

Throughout the paper, let Y be a linear topological space partially ordered by a proper closed convex pointed (i.e., $C \cap (-C) = \{0\}$) cone C . The dual cone to C is denoted by $C^* := \{y^* \in Y^* | \forall y \in C : y^*(y) \geq 0\}$ and the quasi-interior of C^* is defined by $C^\# := \{y^* \in C^* | \forall y \in C \setminus \{0\} : y^*(y) > 0\}$. Furthermore, let X be a linear space, $F : X \rightrightarrows Y$ (with the “ \rightrightarrows ”-notation we denote that F is a set-valued objective function whose function values are sets in Y), and \mathcal{X} a subset of X . As usual, we denote the graph of the set-valued map F by $\text{graph } F := \{(x, y) \in X \times Y \mid y \in F(x)\}$. Furthermore, we define $F(\mathcal{X}) := \cup_{x \in \mathcal{X}} F(x)$.

In set optimization, the following set-relations play an important role, see Young [33], Nishnianidze [24], Kuroiwa [17, 20] and Eichfelder, Jahn [10]. We will use these set-relations to introduce several concepts of robustness.

Definition 1 (Set less order relation ([33, 24, 10])). *Let $C \subset Y$ be a proper closed convex and pointed cone. Furthermore, let $A, B \subset Y$ be arbitrarily chosen sets. Then the set less order relation is defined by*

$$A \preceq_C^s B \iff A \subseteq B - C \text{ and } A + C \supseteq B.$$

Remark 1. *Of course, we have*

$$A \subseteq B - C \iff \forall a \in A \exists b \in B : a \leq_C b$$

and

$$A + C \supseteq B \iff (\forall b \in B \exists a \in A : a \leq_C b).$$

Definition 2 (Upper-type set-relation ([17, 20])). *Let $A, B \subset Y$ be arbitrary chosen sets and $C \subset Y$ a proper closed convex and pointed cone. Then the u-type set-relation \preceq_C^u is defined by*

$$A \preceq_C^u B \iff A \subseteq B - C \iff \forall a \in A \exists b \in B : a \leq_C b.$$

Another important set order relation is the lower-type set-relation:

Definition 3 (Lower-type set-relation ([17, 20])). *Let $A, B \subset Y$ be arbitrarily chosen sets and $C \subset Y$ a proper closed convex and pointed cone. Then the l-type set-relation \preceq_C^l is defined by*

$$A \preceq_C^l B \iff A + C \supseteq B \iff \forall b \in B \exists a \in A : a \leq_C b.$$

Remark 2. *Note that*

(i) $A \subset B - \text{int } C$ and

(ii) $A + N \subset B - C$ for some neighborhood N of the zero vector 0_Y in Y .

are not equivalent when A is not compact. Clearly (ii) implies (i). From a theoretical viewpoint, (ii) may, in some cases, be more appropriate.

Taking into account this property we suppose in Section 3 that the set-valued map $f_{\mathcal{U}}$ in the formulation of the concepts of robustness for multi-objective optimization problems is compact-valued. This is important in the case where we are dealing with $\text{int } C$ in the definition of robustness.

Remark 3. *There is the following relationship between the l-type set-relation \preceq_C^l and the u-type set-relation \preceq_{-C}^u :*

$$A \preceq_C^l B \iff A + C \supseteq B \iff B \subseteq A - (-C) \iff B \preceq_{-C}^u A.$$

To conclude the notation, we introduce a set-valued optimization problem: Consider $F : X \rightrightarrows Y$, and \mathcal{X} a subset of X . Furthermore, let \preceq be a preorder on Y given by Definition 1, 2, 3, respectively. Then a set-valued optimization problem ($\mathcal{SP}-\preceq$) is given by

$$\preceq - \text{minimize } F(x) \text{ subject to } x \in \mathcal{X}, \tag{\mathcal{SP}-\preceq}$$

where minimal solutions of ($\mathcal{SP}-\preceq$) are defined in the following way:

Definition 4 (Minimal solutions of $(\mathcal{SP} - \preceq)$ w.r.t. the preorder \preceq). *Given a set-valued optimization problem $(\mathcal{SP} - \preceq)$, an element $\bar{x} \in \mathcal{X}$ is called a minimal solution to $(\mathcal{SP} - \preceq)$ if*

$$(F(x) \preceq F(\bar{x}) \text{ for some } x \in \mathcal{X}) \implies F(\bar{x}) \preceq F(x).$$

Remark 4. *If we use the set-relation \preceq_C^l introduced in Definition 3 in the formulation of the solution concept, i.e., we study the set-valued optimization problem of $(\mathcal{SP} - \preceq_C^l)$, we observe that this solution concept is based on comparisons among sets of minimal points of values of F . Furthermore, considering the u-type set-relation \preceq_C^u (Definition 2), i.e., considering the problem $(\mathcal{SP} - \preceq_C^u)$ we recognize that this solution concept is based on comparisons of maximal points of values of F . When $\bar{x} \in \mathcal{X}$ is a minimal solution of problem $(\mathcal{SP} - \preceq_C^l)$ there does not exist $x \in S$ such that $F(x)$ is strictly smaller than $F(\bar{x})$ with respect to the set order \preceq_C^l .*

Furthermore, the following definition of a minimizer of a set-valued optimization problem is very often used in the theory of set optimization and given below. However, the solution concept introduced in Definition 4 is more natural and useful as we can see in Example 1.

In the next definition we use the set of minimal elements of a nonempty subset A of Y with respect to C :

$$\text{Min}(A, C) := \{y \in A \mid A \cap (y - (C \setminus \{0\})) = \emptyset\}.$$

Definition 5 (Minimizer of a set-valued optimization problem). *Let $\bar{x} \in \mathcal{X}$ and $(\bar{x}, \bar{y}) \in \text{graph } F$. The pair $(\bar{x}, \bar{y}) \in \text{graph } F$ is called a minimizer of $F : X \rightrightarrows Y$ over \mathcal{X} with respect to C if $\bar{y} \in \text{Min}(F(\mathcal{X}), C)$.*

For our approach to robustness of uncertain multi-objective optimization problems, minimal solutions in the sense of Definition 4 are useful and therefore, when considering robustness concepts, we will deal with this solution concept in the following.

In order to get an insight to the issue of set-valued optimization problems, we give two examples (see Kuroiwa [19]) of set-valued optimization problems.

Example 1. *Consider the set-valued optimization problem*

$$\preceq_C^l \text{-minimize } F_1(x), \quad \text{subject to } x \in \mathcal{X}, \quad (\mathcal{SP} - \preceq_C^l)$$

with $X = \mathbb{R}$, $Y = \mathbb{R}^2$, $C = \mathbb{R}_+^2$, $\mathcal{X} = [0, 1]$ and $F_1 : \mathcal{X} \rightrightarrows Y$ is given by

$$F_1(x) := \begin{cases} [(1, 0), (0, 1)] & \text{if } x = 0 \\ [(1-x, x), (1, 1)] & \text{if } x \in (0, 1], \end{cases}$$

where $[(a, b), (c, d)]$ is the line segment between (a, b) and (c, d) . Only the element $\bar{x} = 0$ is a minimal solution of $(\mathcal{SP} - \preceq_C^l)$. However, all elements $(\bar{x}, \bar{y}) \in \text{graph } F_1$ with $\bar{x} \in [0, 1]$, $\bar{y} = (1 - \bar{x}, \bar{x})$ for $\bar{x} \in (0, 1]$ and $\bar{y} = (1, 0)$ for $\bar{x} = 0$ are minimizers of the set-valued optimization problem in the sense of Definition 5. This example shows that the solution concept with respect to the set-relation \preceq_C^l (see Definitions 3 and 4) is more natural and useful than the concept of minimizers introduced in Definition 5.

Example 2. *In this example we are looking for minimal solutions of a set-valued optimization problem with respect to the set-relation \preceq_C^u introduced in Definition 2.*

$$\preceq_C^u \text{-minimize } F_2(x), \quad \text{subject to } x \in \mathcal{X}, \quad (\mathcal{SP} - \preceq_C^u)$$

with $X = \mathbb{R}$, $Y = \mathbb{R}^2$, $C = \mathbb{R}_+^2$, $\mathcal{X} = [0, 1]$ and $F_2 : \mathcal{X} \rightrightarrows Y$ is given by

$$F_2(x) := \begin{cases} [[(1, 1), (2, 2)]] & \text{if } x = 0 \\ [[(0, 0), (3, 3)]] & \text{if } x \in (0, 1], \end{cases}$$

where $[[a, b], [c, d]] := \{(y_1, y_2) \mid a \leq y_1 \leq c, b \leq y_2 \leq d\}$. Then the only minimal solution of $(\mathcal{SP} - \preceq_C^u)$ in the sense of Definition 4 is $\bar{x} = 0$.

A visualization of both above discussed examples is given in Figure 1.

In Section 3, we will apply the preorders introduced in Definitions 1, 2, 3 in order to define several concepts of robustness for uncertain multi-objective optimization problems.




Figure 1: Feasible solution sets of F_1 , F_2 , described in Example 1 and 2.

3 Concepts of robustness for multi-objective optimization problems based on set relations and corresponding algorithms

Talking about an uncertain optimization problem, we consider the uncertain data to be given as a parameter (also called *scenario*) $\xi \in \mathcal{U}$ where $\mathcal{U} \subseteq \mathbb{R}^m$ is the so called *uncertainty set*. For each realization of this parameter $\xi \in \mathcal{U}$ we obtain a single optimization problem

$$\begin{aligned} f(x, \xi) &\rightarrow \min \\ \text{s.t. } x &\in \mathcal{X}, \end{aligned} \quad (\mathcal{P}(\xi))$$

where $f : X \times \mathcal{U} \mapsto Y$ is the objective function and $\mathcal{X} \subseteq X$ is the set of feasible solutions (note that we assume the feasible set to be unchanged for every realization of the uncertain parameter). We use the notation

$$f_{\mathcal{U}}(x) := \{f(x, \xi) | \xi \in \mathcal{U}\} \quad (1)$$

for the image of the uncertainty set \mathcal{U} and x under f (note that $f_{\mathcal{U}}(x)$ in general is a set and not a singleton).

Taking into account the discussion in Remark 2 we assume that the set-valued map $f_{\mathcal{U}}$ is compact-valued.

Now, when searching for an optimal solution, one has to overcome the problem that we do not know anything about the different scenarios, e.g., which one is most likely to occur, any kind of probability distribution and so on. Therefore, an uncertain (multi-objective) optimization problem is defined as the family of optimization problems

$$(\mathcal{P}(\xi), \xi \in \mathcal{U}). \quad (\mathcal{P}(\mathcal{U}))$$

Now it is not clear what solution to this problem $(\mathcal{P}(\mathcal{U}))$ would be seen as desirable. Throughout the paper we discuss several concepts of robustness and derive new approaches to robustness for multi-objective optimization problems.

In this section we extend the robustness concepts presented in [8] to general spaces using the preorders introduced in Definitions 1, 2, 3. In particular, we are interested in extending the theorems which provide the foundation for the algorithms for calculating the respective robust solutions. We shortly repeat the various concepts which relate to different set orderings, extend the theorems and then formulate the algorithms. With this, we present some ideas for solving special set-valued optimization problems in this section.

3.1 \preceq_C^u -robustness

We extend the definitions and results presented by Ehrgott et al. [8] about minmax robust efficiency.

Here, a feasible solution $x \in \mathcal{X}$ to $(\mathcal{P}(\mathcal{U}))$ is called *minmax robust efficient* if there is no other feasible solution $\bar{x} \in \mathcal{X} \setminus \{x\}$, such that

$$f_{\mathcal{U}}(\bar{x}) \subseteq f_{\mathcal{U}}(x) - \mathbb{R}_{\geq}^k$$

where $\mathbb{R}_{\geq}^k := \{\lambda \in \mathbb{R}^k : \lambda_i \geq 0 \forall i = 1, \dots, k\}$,

With the definitions of *upper-type set-relation*, see Definition 2, and minmax robust efficiency in mind we can see the close connection between minmax robust efficiency and the upper-type set-relation, since a

solution $x \in \mathcal{X}$ to $(\mathcal{P}(\mathcal{U}))$ is minmax robust efficient if there is no other feasible solution $\bar{x} \in \mathcal{X} \setminus \{x\}$, such that

$$f_{\mathcal{U}}(\bar{x}) \preceq_C^u f_{\mathcal{U}}(x),$$

where $Y = \mathbb{R}^k$ and $C = \mathbb{R}_{\geq}^k$.

Since all the concepts considered in this paper are closely related to a set order relation \preceq , in order to keep the names of the concepts readable we call the respective solution \preceq -robust.

In the following definition we use a preorder \preceq_Q^u like in Definition 2 with $Q = C$, $Q = C \setminus \{0\}$ and $Q = \text{int } C$, respectively, instead of \preceq_C^u :

$$A \preceq_Q^u B : \iff A \subseteq B - Q,$$

where $A, B \subset Y$ are arbitrarily chosen sets. If we are dealing with $Q = \text{int } C$ we suppose $\text{int } C \neq \emptyset$.

Using this notation, the concept of minmax robust efficiency can be re-defined as a concept of robustness in the sense of set optimization in the following way:

Definition 6. *Given an uncertain multi-objective optimization problem $(\mathcal{P}(\mathcal{U}))$, a solution $x^0 \in \mathcal{X}$ is called \preceq_Q^u -robust for $(\mathcal{P}(\mathcal{U}))$ with $Q = C$, $Q = C \setminus \{0\}$ and $Q = \text{int } C$, respectively, if there is no solution $\bar{x} \in \mathcal{X} \setminus \{x^0\}$ such that*

$$f_{\mathcal{U}}(\bar{x}) \preceq_Q^u f_{\mathcal{U}}(x^0).$$

Note that the definition of \preceq_Q^u -robustness is valid for general spaces and general cones C , while the definition of minmax robust efficiency in [8] is for $Y = \mathbb{R}^k$ and $C = \mathbb{R}_{\geq}^k$ only.

The motivation behind this concept is the following: When comparing sets with the u -type set-relation, the upper bounds of these sets, i.e., the “worst cases”, are considered. Minimizing these worst cases is closely connected to the concept of minmax robust efficiency where one wants to minimize the objective function in the worst case. This risk averse approach would reflect a decision makers strategy to hedge against a worst case and is rather pessimistic.

Remark 5. *The first scenario-based concept to uncertain multi-objective optimization, or minmax-robustness adapted to multi-objective optimization, has been introduced by Kuroiwa and Lee [21] and studied in [22]. In [21, 22] robust solutions of multi-objective optimization problems are introduced in the following way. The authors propose to consider the robust counterpart to $(\mathcal{P}(\mathcal{U}))$*

$$\text{Min}(f_{RC}^{\mathcal{U}}(\mathcal{X}), \mathbb{R}_{\geq}^k), \quad (2)$$

where the objective vector for $x \in \mathcal{X}$ is given by

$$f_{RC}^{\mathcal{U}}(x) := \begin{pmatrix} \max_{\xi_1 \in \mathcal{U}_1} f_1(x, \xi_1) \\ \dots \\ \max_{\xi_k \in \mathcal{U}_k} f_k(x, \xi_k) \end{pmatrix}, \quad (3)$$

with functionals $f_i : \mathbb{R}^n \times \mathcal{U}_i \rightarrow \mathbb{R}$ for $i = 1, \dots, k$ and the convex and compact uncertainty sets $\mathcal{U} := (\mathcal{U}_1, \dots, \mathcal{U}_k)$ ($\mathcal{U}_i \subseteq \mathbb{R}^m$, $i = 1, \dots, k$). In [21], solutions to (2) are called robust.

Note that in [8] the authors pointed out that this concept differs from the concept of minmax robust efficiency.

With the definition of \preceq_C^u -robustness, we can generalize algorithms for computing minmax robust efficient solutions which is an extension of the well-known weighted sum scalarization technique for calculating efficient solutions to multi-objective optimization problems (compare e.g. Ehrgott [9]).

The general idea is to form a scalar optimization problem by multiplying each objective function with a positive weight and summing up the weighted objectives. The resulting (single-objective) problem in a more general setting is

$$\min_{x \in \mathcal{X}} \sup_{\xi \in \mathcal{U}} y^* \circ f(x, \xi), \quad (\mathcal{P}(\mathcal{U})_{y^*})$$

where $f : X \times \mathcal{U} \rightarrow Y$ and $y^* \in C^* \setminus \{0\}$, i.e., $y^* : Y \rightarrow \mathbb{R}$.

Now, solving this problem one can obtain \preceq_C^u -robust solutions as shown in Theorem 4.3 in [8]. Before extending this theorem, we need a lemma which will help during the proofs:

Lemma 1. Consider the uncertain multi-objective optimization problem $(\mathcal{P}(\mathcal{U}))$. Then it holds for all $x', \bar{x} \in \mathcal{X}$ and for $Q = \text{int } C$ ($Q = C \setminus \{0\}$, $Q = C$, respectively),

$$f_{\mathcal{U}}(x') \subseteq f_{\mathcal{U}}(\bar{x}) - Q \iff \forall \xi \in \mathcal{U} \exists \eta \in \mathcal{U} : f(x', \xi) \in f(\bar{x}, \eta) - Q. \quad (4)$$

Proof. " \implies ": Suppose the contrary. Then

$$\begin{aligned} \exists \xi \in \mathcal{U} \forall \eta \in \mathcal{U} : f(x', \xi) \notin f(\bar{x}, \eta) - Q &\implies \exists \xi \in \mathcal{U} : f(x', \xi) \notin f_{\mathcal{U}}(\bar{x}) - Q \\ &\implies f_{\mathcal{U}}(x') \not\subseteq f_{\mathcal{U}}(\bar{x}) - Q. \end{aligned}$$

" \impliedby ": Suppose the contrary. Then

$$\exists \xi \in \mathcal{U} : f(x', \xi) \notin f_{\mathcal{U}}(\bar{x}) - Q \implies \exists \xi \in \mathcal{U} \forall \eta \in \mathcal{U} : f(x', \xi) \notin f(\bar{x}, \eta) - Q.$$

□

With this, we can extend Theorem 4.3 from [8] in the following way:

Theorem 1. Consider an uncertain multi-objective optimization problem $(\mathcal{P}(\mathcal{U}))$. The following statements hold:

- (a) If $x^0 \in \mathcal{X}$ is a unique optimal solution of $(\mathcal{P}(\mathcal{U})_{y^*})$ for some $y^* \in C^* \setminus \{0\}$, then x^0 is a \preceq_C^u -robust solution for $(\mathcal{P}(\mathcal{U}))$.
- (b) If $x^0 \in \mathcal{X}$ is an optimal solution of $(\mathcal{P}(\mathcal{U})_{y^*})$ for some $y^* \in C^\#$ and $\max_{\xi \in \mathcal{U}} y^* \circ f(x, \xi)$ exists for all $x \in \mathcal{X}$, then x^0 is a $\preceq_{C \setminus \{0\}}^u$ -robust solution for $(\mathcal{P}(\mathcal{U}))$.
- (c) If $x^0 \in \mathcal{X}$ is an optimal solution of $(\mathcal{P}(\mathcal{U})_{y^*})$ for some $y^* \in C^* \setminus \{0\}$ and $\max_{\xi \in \mathcal{U}} y^* \circ f(x, \xi)$ exists for all $x \in \mathcal{X}$, then x^0 is a $\preceq_{\text{int } C}^u$ -robust solution for $(\mathcal{P}(\mathcal{U}))$.

Proof. Suppose that x^0 is not \preceq_Q^u -robust for $Q = C$, $Q = (C \setminus \{0\})$, $Q = \text{int } C$, respectively. Then there exists an element $\bar{x} \in \mathcal{X} \setminus \{x^0\}$ such that

$$f_{\mathcal{U}}(\bar{x}) \subseteq f_{\mathcal{U}}(x^0) - Q, \quad (5)$$

for $Q = C$, ($Q = (C \setminus \{0\})$, $Q = \text{int } C$, respectively).

This implies

$$\forall \xi \in \mathcal{U} \exists \eta \in \mathcal{U} : f(\bar{x}, \xi) \in f(x^0, \eta) - Q.$$

taking into account Lemma 1.

Choose now $y^* \in C^* \setminus \{0\}$ for $Q = C$ ($y^* \in C^\#$ for $Q = C \setminus \{0\}$, $y^* \in (C^* \setminus \{0\})$ for $Q = \text{int } C$, respectively) arbitrary but fixed.

$$\begin{aligned} &\implies \forall \xi \in \mathcal{U} \exists \eta \in \mathcal{U} : y^* \circ f(\bar{x}, \xi) \leq (<, <, \text{ respectively}) y^* \circ f(x^0, \eta) \\ &\implies \forall \xi \in \mathcal{U} : y^* \circ f(\bar{x}, \xi) \leq (<, <, \text{ respectively}) \sup_{\eta' \in \mathcal{U}} y^* \circ f(x^0, \eta') \\ &\implies \sup_{\xi' \in \mathcal{U}} y^* \circ f(\bar{x}, \xi') \leq (<, <, \text{ respectively}) \sup_{\eta' \in \mathcal{U}} y^* \circ f(x^0, \eta'). \end{aligned}$$

The last equation holds because for (b) and (c) $\max_{\xi' \in \mathcal{U}} y^* \circ f(\bar{x}, \xi')$ exists. But this means that x^0 is not the unique optimal (an optimal, an optimal, respectively) solution of $(\mathcal{P}(\mathcal{U})_{y^*})$ for $y^* \in C^* \setminus \{0\}$ ($y^* \in C^\#$, $y^* \in C^* \setminus \{0\}$, respectively). □

Remark 6. In Theorem 1 (b) we consider $y^* \in C^\#$. Under our assumptions concerning the cone C and if we assume additionally $Y = \mathbb{R}^q$ we have $C^\# \neq \emptyset$ (compare [13, Theorem 2.2.12], [13, Example 2.2.16]). Moreover, if Y is a Hausdorff locally convex space, $C \subset Y$ is a proper convex cone and C has a base B with $0 \notin \text{cl } B$, then $C^\# \neq \emptyset$ (compare [13, Theorem 2.2.12]).

With this theorem we can now formulate a first algorithm for finding \preceq_Q^u -robust solutions for $Q = C$, $Q = C \setminus \{0\}$, $Q = \text{int } C$, respectively.

Algorithm 1 for deriving $(\preceq_C^u, \preceq_{C \setminus \{0\}}^u, \preceq_{\text{int } C}^u)$ -robust solutions to $(\mathcal{P}(\mathcal{U}))$ based on weighted sum scalarization:

Input: Uncertain multi-objective problem $\mathcal{P}(\mathcal{U})$, solution sets $\text{Opt}_C = \text{Opt}_{C \setminus \{0\}} = \text{Opt}_{\text{int } C} = \emptyset$.

Step 1: Choose a set $\bar{C} \subset C^* \setminus \{0\}$.

Step 2: If $\bar{C} = \emptyset$: **STOP. Output:** Set of \preceq_C^u -robust solutions Opt_C , set of $\preceq_{C \setminus \{0\}}^u$ -robust solutions $\text{Opt}_{C \setminus \{0\}}$, set of $\preceq_{\text{int } C}^u$ -robust solutions $\text{Opt}_{\text{int } C}$.

Step 3: Choose $y^* \in \bar{C}$. Set $\bar{C} := \bar{C} \setminus \{y^*\}$.

Step 4: Find an optimal solution x^0 of $(\mathcal{P}(\mathcal{U})_{y^*})$.

a) If x^0 is a unique optimal solution of $(\mathcal{P}(\mathcal{U})_{y^*})$, then x^0 is \preceq_C^u -robust for $(\mathcal{P}(\mathcal{U}))$, thus

$$\text{Opt}_C := \text{Opt}_C \cup \{x^0\}.$$

b) If $\max_{\xi \in \mathcal{U}} y^* \circ f(x, \xi)$ exists for all $x \in \mathcal{X}$, then x^0 is $\preceq_{\text{int } C}^u$ -robust for $(\mathcal{P}(\mathcal{U}))$, thus

$$\text{Opt}_{\text{int } C} := \text{Opt}_{\text{int } C} \cup \{x^0\}.$$

c) If $\max_{\xi \in \mathcal{U}} y^* \circ f(x, \xi)$ exists for all $x \in \mathcal{X}$ and $y^* \in C^\#$, then x^0 is $\preceq_{C \setminus \{0\}}^u$ -robust for $(\mathcal{P}(\mathcal{U}))$, thus

$$\text{Opt}_{C \setminus \{0\}} := \text{Opt}_{C \setminus \{0\}} \cup \{x^0\}.$$

Step 5: Go to Step 2.

Furthermore, we present an interactive algorithm for finding $(\preceq_C^u, \preceq_{C \setminus \{0\}}^u, \preceq_{\text{int } C}^u)$ -robust solutions to the uncertain multi-objective optimization problem $(\mathcal{P}(\mathcal{U}))$. This algorithm uses the input of the decision maker whether she accepts the calculated solution or not:

Algorithm 2 for deriving a single accepted $(\preceq_C^u, \preceq_{C \setminus \{0\}}^u, \preceq_{\text{int } C}^u)$ -robust solution to $(\mathcal{P}(\mathcal{U}))$ based on weighted sum scalarization:

Input: Uncertain multi-objective problem $(\mathcal{P}(\mathcal{U}))$.

Step 1: Choose a set $\bar{C} \subset C^* \setminus \{0\}$.

Step 2: Set $j := 0$, choose $\bar{y}^j \in \bar{C}$. Set $\bar{C} := \bar{C} \setminus \{\bar{y}^j\}$.

Step 3: Find an optimal solution x^0 to $(\mathcal{P}(\mathcal{U})_{\bar{y}^j})$.

a) If x^0 is a unique optimal solution of $(\mathcal{P}(\mathcal{U})_{\bar{y}^j})$, then x^0 is \preceq_C^u -robust for $(\mathcal{P}(\mathcal{U}))$.

b) If $\max_{\xi \in \mathcal{U}} \bar{y}^j \circ f(x, \xi)$ exists for all $x \in \mathcal{X}$, then x^0 is $\preceq_{\text{int } C}^u$ -robust for $(\mathcal{P}(\mathcal{U}))$.

c) If $\max_{\xi \in \mathcal{U}} \bar{y}^j \circ f(x, \xi)$ exists for all $x \in \mathcal{X}$ and $\bar{y}^j \in C^\#$, then x^0 is $\preceq_{C \setminus \{0\}}^u$ -robust for $(\mathcal{P}(\mathcal{U}))$.

If x^0 is accepted by the decision-maker: **STOP. Output:** x^0 .

Step 4: Choose $\bar{y}^{j+1} \in C$, such that $\bar{y}^{j+1} \neq \bar{y}^i$ for all $0 \leq i \leq j$.

Set $l := 0$, $t_0 := 0$.

Step 5: Choose t_{l+1} with $t_l < t_{l+1} \leq 1$ and compute an optimal solution x_{l+1}^j to

$$\min_{x \in \mathcal{X}} \sup_{\xi \in \mathcal{U}} \bar{y}^* \circ f(x, \xi) \quad (\mathcal{P}(\mathcal{U})_{\bar{y}^j + t_{l+1}(\bar{y}^{j+1} - \bar{y}^j)})$$

with $\bar{y}^* := \bar{y}^j + t_{l+1}(\bar{y}^{j+1} - \bar{y}^j)$. If no optimal solution to $(\mathcal{P}(\mathcal{U})_{\bar{y}^j + t_{l+1}(\bar{y}^{j+1} - \bar{y}^j)})$ can be found for all $t > t^l$, go to Step 2.

Step 6: If x_{l+1}^j is accepted by the decision maker: **STOP. Output:** x_{l+1}^j .

Step 7: If $t_{l+1} = 1$, then set $j := j + 1$ and go to Step 4. Otherwise, set $l := l + 1$ and go to Step 5.

3.2 \preceq_C^l -robustness

In this section we use the l -type set-relation \preceq_Q^l like in Definition 3 with $Q = C$, $Q = C \setminus \{0\}$ and $Q = \text{int } C$, respectively, instead of \preceq_C^l :

$$A \preceq_Q^l B : \iff A + Q \supseteq B,$$

where $A, B \subset Y$ are arbitrarily chosen sets. If we are dealing with $Q = \text{int } C$ we suppose $\text{int } C \neq \emptyset$. Using this notation we derive the new concept of \preceq_Q^l -robustness, defined analogously to \preceq_Q^u -robustness (Definition 6):

Definition 7. *Given an uncertain multi-objective optimization problem $(\mathcal{P}(\mathcal{U}))$, a solution $x^0 \in \mathcal{X}$ is called \preceq_Q^l -robust if there is no $\bar{x} \in \mathcal{X} \setminus \{x^0\}$ such that*

$$f_{\mathcal{U}}(\bar{x}) \preceq_Q^l f_{\mathcal{U}}(x^0).$$

The \preceq_Q^l -robustness (with $Q = C$, $Q = C \setminus \{0\}$ and $Q = \text{int } C$, respectively) can be interpreted as an optimistic approach. The following example illustrates this concept for the case $Q = C$.

Remark 7. *Here, x is \preceq_C^l -robust, while it is not \preceq_C^u -robust.*




Figure 2: x is \preceq_C^l -robust.

The \preceq_Q^l -robustness is an alternative tool for the decision maker for obtaining solutions of another type to an uncertain multi-objective optimization problem. This rather optimistic approach focuses on the lower bound of a set $f_{\mathcal{U}}(\bar{x})$ for the comparison to another set $f_{\mathcal{U}}(x^0)$. In particular, in the case $Q = C$, a point $x^0 \in \mathcal{X}$ is called a \preceq_C^l -solution if there is no other point $\bar{x} \in \mathcal{X}$ such that $f_{\mathcal{U}}(x^0)$ is a subset of $f_{\mathcal{U}}(\bar{x}) + C$. Contrary to the \preceq_Q^u -robustness approach, the \preceq_Q^l -robustness (with $Q = C$, $Q = C \setminus \{0\}$ and $Q = \text{int } C$, respectively) is hence not a worst-case concept, thus the decision maker is not considered to be risk averse but risk affine. This optimistic concept thus hedges against perturbations in the best-case scenarios.

For calculating \preceq_Q^l -robust solutions again the weighted sum scalarization is helpful, but in order to later on compute \preceq_Q^l -robust solutions to $(\mathcal{P}(\mathcal{U}))$, we define a new weighted sum problem in a general setting:

Let $y^* \in C^* \setminus \{0\}$ ($y \in C^\#$, respectively). Consider the weighted sum scalarization problem

$$\min_{x \in \mathcal{X}} \inf_{\xi \in \mathcal{U}} y^* \circ f(x, \xi). \quad (\mathcal{P}(\mathcal{U})_{y^*}^{opt})$$

Theorem 2. *Consider an uncertain vector optimization problem $(\mathcal{P}(\mathcal{U}))$. The following statements hold:*

- (a) *If x^0 is a unique optimal solution of $(\mathcal{P}(\mathcal{U})_{y^*}^{opt})$ for some $y^* \in C^* \setminus \{0\}$, then x^0 is a \preceq_C^l -robust solution to $(\mathcal{P}(\mathcal{U}))$.*
- (b) *If x^0 is an optimal solution of $(\mathcal{P}(\mathcal{U})_{y^*}^{opt})$ for some $y^* \in C^\#$ and $\min_{\xi \in \mathcal{U}} y^* \circ f(x, \xi)$ exists for all $x \in \mathcal{X}$, then x^0 is a $\preceq_{C \setminus \{0\}}^l$ -robust solution to $(\mathcal{P}(\mathcal{U}))$.*
- (c) *If x^0 is an optimal solution of $(\mathcal{P}(\mathcal{U})_{y^*}^{opt})$ for some $y^* \in C^* \setminus \{0\}$ and $\min_{\xi \in \mathcal{U}} y^* \circ f(x, \xi)$ exists for all $x \in \mathcal{X}$, then x^0 is a $\preceq_{\text{int } C}^l$ -robust solution to $(\mathcal{P}(\mathcal{U}))$.*

Proof. Suppose x^0 is not \preceq_Q^l -robust for $Q = C$ ($Q = C \setminus \{0\}$, $Q = \text{int } C$, respectively). Consequently, there exists an $\bar{x} \in \mathcal{X} \setminus \{x^0\}$ s.t. $f_{\mathcal{U}}(\bar{x}) + Q \supseteq f_{\mathcal{U}}(x^0)$ for $Q = C$ ($Q = C \setminus \{0\}$, $Q = \text{int } C$, respectively). That is equivalent to

$$\begin{aligned} & \forall \xi \in \mathcal{U} \exists \eta \in \mathcal{U} : f(\bar{x}, \eta) + Q \ni f(x^0, \xi) \\ & \iff \forall \xi \in \mathcal{U} \exists \eta \in \mathcal{U} : f(\bar{x}, \eta) \in f(x^0, \xi) - Q. \end{aligned} \quad (6)$$

Now choose $y^* \in C^* \setminus \{0\}$ for $Q = C$ ($y^* \in C^\#$ for $Q = C \setminus \{0\}$, $y^* \in C^* \setminus \{0\}$ for $Q = \text{int } C$, respectively) arbitrary, but fixed. Hence, we obtain from (6)

$$\begin{aligned} & \implies \forall \xi \in \mathcal{U} \exists \eta \in \mathcal{U} : y^* \circ f(\bar{x}, \eta) \leq (<, <, \text{respectively}) y^* \circ f(x^0, \xi) \\ & \implies \exists \eta \in \mathcal{U} : y^* \circ f(\bar{x}, \eta) \leq (<, <, \text{respectively}) \inf_{\xi \in \mathcal{U}} y^* \circ f(x^0, \xi) \\ & \implies \inf_{\eta \in \mathcal{U}} y^* \circ f(\bar{x}, \eta) \leq (<, <, \text{respectively}) \inf_{\xi \in \mathcal{U}} y^* \circ f(x^0, \xi), \end{aligned}$$

in contradiction to the assumption. \square

Based on these results, we are able to present the following algorithm that computes $(\preceq_C^l / \preceq_{C \setminus \{0\}}^l / \preceq_{\text{int } C}^l)$ -robust solutions to $\mathcal{P}(\mathcal{U})$:

Algorithm 3 for deriving $(\preceq_C^l / \preceq_{C \setminus \{0\}}^l / \preceq_{\text{int } C}^l)$ -robust solutions to $(\mathcal{P}(\mathcal{U}))$ based on weighted sum scalarization:

Input & Step 1-5: Analogous to Algorithm 1, only replacing $(\mathcal{P}(\mathcal{U})_{y^*})$ by $(\mathcal{P}(\mathcal{U})_{y^*}^{opt})$ and replacing $\max_{\xi \in \mathcal{U}} y^* \circ f(x^0, \xi)$ by $\min_{\xi \in \mathcal{U}} y^* \circ f(x^0, \xi)$.

The next algorithm computes $(\preceq_C^l / \preceq_{C \setminus \{0\}}^l / \preceq_{\text{int } C}^l)$ -robust solutions via weighted sum scalarization by altering the weights:

Algorithm 4 for calculating a single desired $(\preceq_C^l / \preceq_{C \setminus \{0\}}^l / \preceq_{\text{int } C}^l)$ -robust solution to $(\mathcal{P}(\mathcal{U}))$ based on weighted sum scalarization:

Input & Step 1-7: Analogous to Algorithm 2, only replacing $(\mathcal{P}(\mathcal{U})_{y^*})$ by $(\mathcal{P}(\mathcal{U})_{y^*}^{opt})$, $\max_{\xi \in \mathcal{U}} y^* \circ f(x^0, \xi)$ by $\min_{\xi \in \mathcal{U}} y^* \circ f(x^0, \xi)$ and $(\mathcal{P}(\mathcal{U})_{\bar{y}^j + t_{l+1}(\bar{y}^{j+1} - \bar{y}^j)})$ by $(\mathcal{P}(\mathcal{U})_{\bar{y}^j + t_{l+1}(\bar{y}^{j+1} - \bar{y}^j)}^{opt})$.

3.3 \preceq_C^s -robustness

Now, we use the set less order relation \preceq_Q^s with $Q = C$, $Q = C \setminus \{0\}$ and $Q = \text{int } C$, respectively (compare Definition 1) for $A, B \subset Y$ arbitrarily chosen sets:

$$A \preceq_Q^s B : \iff A \subseteq B - Q \text{ and } A + Q \supseteq B.$$

If we are dealing with $Q = \text{int } C$ we suppose $\text{int } C \neq \emptyset$.

We can now introduce the concept of \preceq_Q^s -robustness (with $Q = C$, $Q = C \setminus \{0\}$ and $Q = \text{int } C$, respectively):

Definition 8. A solution x^0 of $(\mathcal{P}(\mathcal{U}))$ is called $(\preceq_C^s / \preceq_{C \setminus \{0\}}^s / \preceq_{\text{int } C}^s)$ -robust if there is no $\bar{x} \in \mathcal{X} \setminus \{x^0\}$ such that

$$f_{\mathcal{U}}(\bar{x}) \preceq_Q^s f_{\mathcal{U}}(x^0)$$

for $Q = C$ ($Q = C \setminus \{0\}$, $Q = \text{int } C$, respectively).

Remark 8. Figure 3 shows an element $x \in \mathcal{X}$ that is \preceq_C^s -robust, while it is not $\preceq_{\text{int } C}^u$ -robust.

Remark 9. Note that a \preceq_C^l -robust solution is as well \preceq_C^s -robust by definition. The same holds for a \preceq_C^u -robust solution.




Figure 3: x is \preceq_C^s -robust.

The concept of \preceq_C^s -robustness can be interpreted in the following way: In a situation where it is not clear if one should follow a risk affine or risk averse strategy (e.g., the decision maker is not at hand or wants to get a feeling for the variety of the solutions) this concept might be helpful as it calculates solutions which reflect these different strategies. Therefore, this concept can serve as a pre-selection before deciding a definite strategy.

Computing \preceq_C^s -robust solutions is possible with the help of the following optimization problem:

$$h(x) := \left(\begin{array}{c} \inf_{\xi \in \mathcal{U}} y^* \circ f(x, \xi) \\ \sup_{\xi \in \mathcal{U}} y^* \circ f(x, \xi) \end{array} \right) \rightarrow v - \min_{x \in \mathcal{X}} \quad (\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$$

with $y^* \in C^* \setminus \{0\}$ ($y^* \in C^\#$, respectively). For $(\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$, we use the solution concept of weak Pareto efficiency: An element $x^0 \in \mathcal{X}$ is called weakly Pareto efficient for $(\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$, if

$$h(\mathcal{X}) \cap (h(x^0) - \text{int } \mathbb{R}_{\geq}^2) = \emptyset.$$

Furthermore, a point $x^0 \in \mathcal{X}$ is called strictly Pareto efficient for $(\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$, if

$$h(\mathcal{X} \setminus \{x^0\}) \cap (h(x^0) - \mathbb{R}_{\geq}^2) = \emptyset.$$

We have the following theorem:

Theorem 3. Consider an uncertain multi-objective optimization problem $(\mathcal{P}(\mathcal{U}))$. The following statements hold:

- (a) If x^0 is strictly Pareto efficient for problem $(\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$ for some $y^* \in C^* \setminus \{0\}$, then x^0 is \preceq_C^s -robust.
- (b) If x^0 is weakly Pareto efficient for problem $(\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$ for some $y^* \in C^* \setminus \{0\}$ and $\min_{\xi \in \mathcal{U}} y^* \circ f(x, \xi)$ and $\max_{\xi \in \mathcal{U}} y^* \circ f(x, \xi)$ exist for all $x \in \mathcal{X}$ and the chosen weight $y^* \in C^* \setminus \{0\}$, then x^0 is $\preceq_{\text{int } C}^s$ -robust.
- (c) If x^0 is weakly Pareto efficient for problem $(\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$ for some $y^* \in C^\#$ and $\min_{\xi \in \mathcal{U}} y^* \circ f(x, \xi)$ and $\max_{\xi \in \mathcal{U}} y^* \circ f(x, \xi)$ exist for all $x \in \mathcal{X}$ and the chosen weight $y^* \in C^\#$, then x^0 is $\preceq_{C \setminus \{0\}}^s$ -robust.

Proof. Let x^0 be strictly Pareto efficient (weakly Pareto efficient, weakly Pareto efficient) for problem $(\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$ with some $y^* \in C^* \setminus \{0\}$ ($y^* \in C^* \setminus \{0\}$, $y^* \in C^\#$, respectively), i.e., there is no $\bar{x} \in \mathcal{X} \setminus \{x^0\}$ such that

$$\begin{aligned} \inf_{\xi \in \mathcal{U}} y^* \circ f(\bar{x}, \xi) &\leq (<, <, \text{ respectively}) \inf_{\xi \in \mathcal{U}} y^* \circ f(x^0, \xi) \\ \text{and } \sup_{\xi \in \mathcal{U}} y^* \circ f(\bar{x}, \xi) &\leq (<, <, \text{ respectively}) \sup_{\xi \in \mathcal{U}} y^* \circ f(x^0, \xi). \end{aligned}$$

Now suppose x^0 is not $(\preceq_C^s / \preceq_{\text{int } C}^s / \preceq_{C \setminus \{0\}}^s)$ -robust. Then there exists an $\bar{x} \in \mathcal{X} \setminus \{x^0\}$ such that

$$f_{\mathcal{U}}(\bar{x}) + Q \supseteq f_{\mathcal{U}}(x^0) \text{ and } f_{\mathcal{U}}(\bar{x}) \subseteq f_{\mathcal{U}}(x^0) - Q$$

for $Q = C$ ($Q = \text{int } C$, $Q = C \setminus \{0\}$). That implies

$$\exists \bar{x} \in \mathcal{X} \setminus \{x^0\} : \forall \xi_1, \xi_2 \in \mathcal{U} \exists \eta_1, \eta_2 \in \mathcal{U} : f(\bar{x}, \eta_1) + Q \ni f(x^0, \xi_1) \text{ and } f(\bar{x}, \xi_2) \in f(x^0, \eta_2) - Q \quad (7)$$

for $Q = C$ ($Q = \text{int } C$, $Q = C \setminus \{0\}$). Choose now $y^* \in C^* \setminus \{0\}$ ($y^* \in C^* \setminus \{0\}$, $y^* \in C^\#$) as in problem $(\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$. We obtain from (7)

$$\begin{aligned} & \exists \bar{x} \in \mathcal{X} \setminus \{x^0\} : \forall \xi_1, \xi_2 \in \mathcal{U} \exists \eta_1, \eta_2 \in \mathcal{U} : y^* \circ f(\bar{x}, \eta_1) \leq (<, <, \text{ respectively}) y^* \circ f(x^0, \xi_1) \\ & \quad \text{and } y^* \circ f(\bar{x}, \xi_2) \leq (<, <, \text{ respectively}) y^* \circ f(x^0, \eta_2) \\ & \Rightarrow \inf_{\xi \in \mathcal{U}} y^* \circ f(\bar{x}, \xi) \leq (<, <, \text{ respectively}) \inf_{\xi \in \mathcal{U}} y^* \circ f(x^0, \xi) \\ & \quad \text{and } \sup_{\xi \in \mathcal{U}} y^* \circ f(\bar{x}, \xi) \leq (<, <, \text{ respectively}) \sup_{\xi \in \mathcal{U}} y^* \circ f(x^0, \xi). \end{aligned}$$

The last two strict inequalities hold because the minimum and maximum exist. But this is a contradiction to the assumption. \square

Based on these observations, we can formulate the following algorithm for computing \preceq_C^s -robust solutions to $\mathcal{P}(\mathcal{U})$.

Algorithm 5 for computing $(\preceq_C^s / \preceq_{C \setminus \{0\}}^s / \preceq_{\text{int } C}^s)$ -robust solutions using a family of problems $(\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$:

Input & Step 1-3: Analogous to Algorithm 1.

Step 4: Find a set of weakly Pareto efficient solutions $SOL_{we}(y^*)$ of $(\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$.

Step 5: If $SOL_{we}(y^*) = \emptyset$, then go to Step 2.

Step 6: Choose $\bar{x} \in SOL_{we}(y^*)$. Set $SOL_{we}(y^*) := SOL_{we}(y^*) \setminus \{\bar{x}\}$.

a) If \bar{x} is a strictly Pareto efficient solution of $(\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$, then \bar{x} is \preceq_C^s -robust for $(\mathcal{P}(\mathcal{U}))$, thus

$$\text{Opt}_C := \text{Opt}_C \cup \{\bar{x}\}.$$

b) If \bar{x} is a weakly Pareto efficient solution of $(\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$ and $\max_{\xi \in \mathcal{U}} y^* \circ f(x, \xi)$ and $\min_{\xi \in \mathcal{U}} y^* \circ f(x, \xi)$ exist for all $x \in \mathcal{X}$, then \bar{x} is $\preceq_{\text{int } C}^s$ -robust for $(\mathcal{P}(\mathcal{U}))$, thus

$$\text{Opt}_{\text{int } C} := \text{Opt}_{\text{int } C} \cup \{\bar{x}\}.$$

c) If \bar{x} is weakly Pareto efficient for problem $(\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$ and $y^* \in C^\#$ and $\min_{\xi \in \mathcal{U}} y^* \circ f(x, \xi)$ and $\max_{\xi \in \mathcal{U}} y^* \circ f(x, \xi)$ exist for all $x \in \mathcal{X}$ and the chosen weight $y^* \in C^\#$, then \bar{x} is $\preceq_{C \setminus \{0\}}^s$ -robust for $(\mathcal{P}(\mathcal{U}))$, thus

$$\text{Opt}_{C \setminus \{0\}} := \text{Opt}_{C \setminus \{0\}} \cup \{\bar{x}\}.$$

Step 7: Go to Step 5.

In the following we present an algorithm that computes \preceq_C^s -robust solutions while varying the weights in the vector of objectives of problem $(\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$.

Algorithm 6 for computing $(\preceq_C^s / \preceq_{C \setminus \{0\}}^s / \preceq_{\text{int } C}^s)$ -robust solutions using a family of problems $(\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$:

Input & Step 1-3 & Step 5-7: Analogous to Algorithm 2, only replacing $(\mathcal{P}(\mathcal{U})_{y^*})$ by $(\mathcal{P}(\mathcal{U})_{y^*}^{biobj})$ and $(\mathcal{P}(\mathcal{U})_{\bar{y}^j + t_{l+1}(\bar{y}^{j+1} - \bar{y}^j)})$ by $(\mathcal{P}(\mathcal{U})_{\bar{y}^j + t_{l+1}(\bar{y}^{j+1} - \bar{y}^j)}^{biobj})$.

Step 4: Analogous to **Step 4** of Algorithm 5.

3.4 Alternative Set Less Ordered Robustness

Another way of combining the u - and l -type set-relations is the alternative set less order relation:

Definition 9 (Alternative set less order relation (compare Ide and Köbis [14])). *Let $C \subset Y$ be a proper closed convex and pointed cone. Furthermore, let $A, B \subset Y$ be arbitrarily chosen sets. Then the alternative set less order relation is defined by*

$$A \preceq_C^a B \iff A \preceq_C^u B \text{ or } A \preceq_C^l B.$$

Based on this definition we can now define the concept of \preceq_C^a -robustness for general cones:

Definition 10. *A solution x^0 of $(\mathcal{P}(\mathcal{U}))$ is called $(\preceq_C^a / \preceq_{C \setminus \{0\}}^a / \preceq_{\text{int } C}^a)$ -robust if there is no $\bar{x} \in \mathcal{X} \setminus \{x^0\}$ such that*

$$f_{\mathcal{U}}(\bar{x}) \preceq_Q^a f_{\mathcal{U}}(x^0)$$

for $Q = C$ ($Q = C \setminus \{0\}$, $Q = \text{int } C$, respectively).

The following example illustrates \preceq_C^a -robust solutions.

Remark 10. *In Figure 4, both x and \bar{x} are \preceq_C^a -robust.*




Figure 4: Both x and \bar{x} are strictly \preceq_C^a -robust.

The next lemma follows directly from the definitions:

Lemma 2. *Note that a solution of $(\mathcal{P}(\mathcal{U}))$ is \preceq_C^a -robust if and only if it is \preceq_C^l -robust and \preceq_C^u -robust.*

As this lemma shows, the concept of \preceq_C^a -robustness is rather restrictive as only solutions which are \preceq_C^u -robust and \preceq_C^l -robust, thus reflect both a risk averse and a risk affine strategy, are also \preceq_C^a -robust. Therefore, this concept is fit for a decision maker who does not want to make any mistake in terms of the best or worst cases. We can see easily that such an approach would be very restrictive against the solutions and that only very few solutions should fulfill these conditions.

Due to this Lemma 2, from Algorithms 1 and 3, we can deduce the following algorithm for calculating \preceq_C^a -robust solutions to $(\mathcal{P}(\mathcal{U}))$:

Algorithm 7 for deriving $(\preceq_C^a / \preceq_{C \setminus \{0\}}^a / \preceq_{\text{int } C}^a)$ -robust solutions to $(\mathcal{P}(\mathcal{U}))$:

Input: Uncertain multi-objective problem $(\mathcal{P}(\mathcal{U}))$, solution sets $\text{Opt}_C^a = \text{Opt}_{C \setminus \{0\}}^a = \text{Opt}_{\text{int } C}^a = \emptyset$.

Step 1: Compute a set of $(\preceq_C^l / \preceq_{\text{int } C}^l / \preceq_{C \setminus \{0\}}^l)$ -robust solutions $(\text{Opt}_C^l, \text{Opt}_{\text{int } C}^l, \text{Opt}_{C \setminus \{0\}}^l)$ using Algorithm 3 or 4.

Step 2: Compute a set of $(\preceq_C^u / \preceq_{\text{int } C}^u / \preceq_{C \setminus \{0\}}^u)$ -robust solutions $(\text{Opt}_C^u, \text{Opt}_{\text{int } C}^u, \text{Opt}_{C \setminus \{0\}}^u)$ using Algorithm 1 or 2.

Output: Set of $(\preceq_C^a / \preceq_{\text{int } C}^a / \preceq_{C \setminus \{0\}}^a)$ -robust solutions

$$\begin{aligned} \text{Opt}_C^a &= \text{Opt}_C^u \cap \text{Opt}_C^l, \\ \text{Opt}_{\text{int } C}^a &= \text{Opt}_{\text{int } C}^u \cap \text{Opt}_{\text{int } C}^l, \\ \text{Opt}_{C \setminus \{0\}}^a &= \text{Opt}_{C \setminus \{0\}}^u \cap \text{Opt}_{C \setminus \{0\}}^l. \end{aligned}$$

3.5 Further Relationships Between the Concepts

From Remark 9 we can see that every \preceq_C^u -robust solution and every \preceq_C^l -robust solution is also a \preceq_C^s -robust solution. The inverse direction does not hold. The following example in Figure 5 shows that a solution can be \preceq_C^s -robust but neither \preceq_C^u -robust nor \preceq_C^l -robust.




Figure 5: x is \preceq_C^s -robust, but neither \preceq_C^u -robust nor \preceq_C^l -robust.

We summarize the relationship between the various robustness concepts in Figure 6.




Figure 6: Scheme of solutions to an uncertain multicriteria optimization problem.

4 Conclusions

In the following we will explain that our algorithms presented in Section 3 can be used for solving special classes of set-valued optimization problems.

Having a close look at all the concepts of robustness from Section 3, we can see that in fact all of these are set-valued optimization problems.

Consider a set-valued optimization problem of the form

$$\preceq - \text{minimize } F(x), \quad \text{subject to } x \in \mathcal{X}, \quad (\mathcal{SP} - \preceq)$$

with some given preorder \preceq and a set-valued objective map $F : \mathcal{X} \rightrightarrows Y$, we can see the following:

If the preorder \preceq is given by \preceq_C^a , \preceq_C^l , \preceq_C^u , or \preceq_C^s with some proper closed convex pointed cone $C \subset Y$ and $F(x)$ can be parametrized by parameters $\xi \in \mathcal{U}$ with some set \mathcal{U} in the way that

$$F(x) := f_{\mathcal{U}}(x) \text{ for all } x \in \mathcal{X},$$

where $f_{\mathcal{U}}(x) = \{f(x, \xi) | \xi \in \mathcal{U}\}$ and $f : X \times \mathcal{U} \mapsto Y$, then the set-valued optimization problem $(\mathcal{SP} - \preceq)$ is equivalent to finding \preceq -robust solutions to the uncertain multi-objective problem $(\mathcal{P}(\mathcal{U}))$ and can therefore be solved by using one of the respective algorithms presented in Section 3.

We revealed strong connections between set-valued optimization and uncertain multi-objective optimization. Furthermore, we derived our results in a more general setting than in [8] and [14]. In particular, we provided solution algorithms for a certain class of set-valued optimization problems. It seems possible to extend this class of problems to a more general one, but this is future work and of interest for the next steps in this area of research.

Moreover, this paper made very clear that finding robust solutions to uncertain multi-objective optimization problems can be interpreted as an application of set-valued optimization. Thus, robust solutions to uncertain multi-objective optimization problems can be obtained by using the solution techniques from set-valued optimization. Formulating concrete algorithms of this kind is another topic for future research.

References

- [1] Koji Aoyama, Fumiaki Kohsaka, and Wataru Takahashi. Proximal point methods for monotone operators in Banach spaces. *Taiwanese J. Math.*, 15(1):259–281, 2011.
- [2] C. Barrico and C.H. Antunes. Robustness analysis in multi-objective optimization using a degree of robustness concept. In *IEEE Congress on Evolutionary Computation. CEC 2006.*, pages 1887–1892. IEEE Computer Society, 2006.
- [3] A. Ben-Tal, L. El Ghaoui, and A. Nemirovski. *Robust Optimization*. Princeton University Press, Princeton and Oxford, 2009.
- [4] A. Ben-Tal and A. Nemirovski. Robust convex optimization. *Mathematics of Operations Research*, 23(4):769–805, 1998.
- [5] J.R. Birge and F.V. Louveaux. *Introduction to Stochastic Programming*. Springer Verlag, New York, 1997.
- [6] J. Branke. Creating robust solutions by means of evolutionary algorithms. In E.A. Eiben, T. Bäck, M. Schenauer, and H.-P. Schwefel, editors, *Parallel Problem Solving from Nature – PPSNV*, volume 1498 of *Lecture Notes in Computer Science*, pages 119–128. Springer, Berlin, Heidelberg, 1998.
- [7] K. Deb and H. Gupta. Introducing robustness in multi-objective optimization. *Evolutionary Computation*, 14(4):463–494, 2006.
- [8] M. Ehrgott, J. Ide, and A. Schöbel. Minmax robustness for multi-objective optimization problems.
- [9] Matthias Ehrgott. *Multicriteria optimization*. Springer-Verlag, Berlin, second edition, 2005.
- [10] Gabriele Eichfelder and Johannes Jahn. Vector optimization problems and their solution concepts. In *Recent developments in vector optimization*, Vector Optim., pages 1–27. Springer, Berlin, 2012.
- [11] M. Fischetti and M. Monaci. Light robustness. In R. K. Ahuja, R. Moehring, and C. Zaroliagis, editors, *Robust and Online Large-Scale Optimization*, number 5868 in *Lecture Notes in Computer Science*, pages 61–84. Springer, Berlin Heidelberg, 2009.
- [12] M. Goerigk and A. Schöbel. A scenario-based approach for robust linear optimization. In *Proceedings of the 1st International ICST Conference on Practice and Theory of Algorithms in (Computer) Systems (TAPAS)*, *Lecture Notes in Computer Science*, pages 139–150. Springer, 2011.
- [13] A. Göpfert, H. Riahi, Chr. Tammer, and C. Zălinescu. *Variational Methods in Partially Ordered Spaces*. Springer, New York, 2003.
- [14] J. Ide and E. Köbis. Concepts of robustness for multi-objective optimization problems based on set order relations. 2013.
- [15] K. Klamroth, E. Köbis, A. Schöbel, and Chr. Tammer. A unified approach for different concepts of robustness and stochastic programming via nonlinear scalarizing functionals. *Optimization*, 62:649–671, 2013.
- [16] P. Kouvelis and G. Yu. *Robust Discrete Optimization and its Applications*. Kluwer Academic Publishers, 1997.

- [17] D. Kuroiwa. Some duality theorems of set-valued optimization with natural criteria. In *Proceedings of the International Conference on Nonlinear Analysis and Convex Analysis*. World Scientific, 221–228, 1999.
- [18] Daishi Kuroiwa. On set-valued optimization. *Nonlinear Anal.*, 47(2):1395–1400, 2001.
- [19] Daishi Kuroiwa. Existence theorems of set optimization with set-valued maps. *J. Inf. Optim. Sci.*, 24(1):73–84, 2003.
- [20] Daishi Kuroiwa. The natural criteria in set-valued optimization. *Sūrikaiseikikenkyūsho Kōkyūroku*, (1031):85–90, Research on nonlinear analysis and convex analysis. Kyoto, 1997.
- [21] Daishi Kuroiwa and Gue Myung Lee. On robust multiobjective optimization. *Vietnam Journal of Mathematics*, 40(2&3):305–317, 2012.
- [22] Daishi Kuroiwa and Gue Myung Lee. On robust convex multiobjective optimization. *Accepted to Journal of Nonlinear and Convex Analysis*, 2013.
- [23] C. Liebchen, M. Lübbecke, R. H. Möhring, and S. Stiller. The concept of recoverable robustness, linear programming recovery, and railway applications. In R. K. Ahuja, R.H. Möhring, and C.D. Zaroliagis, editors, *Robust and online large-scale optimization*, volume 5868 of *Lecture Note on Computer Science*. Springer, 2009.
- [24] Z. G. Nishnianidze. Fixed points of monotone multivalued operators. *Soobshch. Akad. Nauk Gruzin. SSR*, 114(3):489–491, 1984.
- [25] E. Marian Scott, Andrea Saltelli, and Tine Sörensen. Practical experience in applying sensitivity and uncertainty analysis. In *Sensitivity analysis*, Wiley Ser. Probab. Stat., pages 267–274. Wiley, Chichester, 2000.
- [26] A.L. Soyster. Convex programming with set-inclusive constraints and applications to inexact linear programming. *Oper. Res.*, 21:1154–1157, 1973.
- [27] Ingrida Steponavičė and Kaisa Miettinen. Survey on multiobjective robustness for simulation-based optimization. *Talk at the 21st International Symposium on Mathematical Programming, August 19-24 2012, Berlin, Germany*, 2012.
- [28] T. Stewart, O. Bandte, H. Braun, N. Chakraborti, M. Ehrgott, M. Göbelt, Y. Jin, H. Nakayama, S. Poles, and D. Di Stefano. Real-world applications of multiobjective optimization. In J. Branke, K. Deb, K. Miettinen, and R. Slowinski, editors, *Multiobjective Optimization: Interactive and Evolutionary Approaches*, volume 5252 of *Lecture Notes in Computer Science*, pages 285–327. Springer, Berlin, 2008.
- [29] Wataru Takahashi. *Nonlinear functional analysis*. Yokohama Publishers, Yokohama, 2000. Fixed point theory and its applications.
- [30] Wataru Takahashi. *Introduction to nonlinear and convex analysis*. Yokohama Publishers, Yokohama, 2009.
- [31] Wataru Takahashi. Nonlinear mappings in equilibrium problems and an open problem in fixed point theory. In *Fixed point theory and its applications*, pages 177–197. Yokohama Publ., Yokohama, 2010.
- [32] K. Witting. *Numerical algorithms for the treatment of parametric multiobjective optimization problems and applications*. PhD thesis, Universität Paderborn, Paderborn, 2012.
- [33] Rosalind Cecily Young. The algebra of many-valued quantities. *Math. Ann.*, 104(1):260–290, 1931.

Reports of the Institutes 2013

- 01-13.** B. Soleimani and Chr. Tammer, *Concepts for approximate solutions of vector optimization problems with variable order structure*
- 02-13.** E. Köbis, *On Robust Optimization: Relations Between Scalar Robust Optimization and Unconstrained Multicriteria Optimization*
- 03-13.** B. Soleimani and Chr. Tammer, *Scalarization of approximate solutions of vector optimization problem with variable order structure based on nonlinear scalarization*